# Speciale



Future scenarios of European forests

Scenarios that are based on contrasting storylines can be used as a tool to explore the different ways in which the future may develop and their impacts on the sustainability of European forestry. These scenarios are neither predictions nor forecasts, but are used to create a consistent image of a future, and to help decision makers and other actors in making well informed choices. Each storyline assumes a distinctly different direction for future developments. Here we discuss characteristics of scenario studies in general, and focus on few recent trends and their long term impact on European forests as, e.g., nature oriented management, carbon credits, and extra demand for bioenergy

■ Gert-Jan Nabuurs

#### Introduction

European forests cover 36% of the European land area, and they fulfil a multitude of functions; from wood production to nature conservation, climate mitigation, water protection, etc. It's this multitude of functions that is achieved by a huge variety of forest owners and national legislations that make these forests unique in the world. Europe namely went through a centuries long period of deforestation and degradation which already started before the Roman Era. Then in Medieval times, wood became a scarce product in many regions, resulting in the first ideas of a sustainably managed forest as published by Von Carlowitz (1713). This has resulted in many new plantings and also after WWII large scale restoration and afforestation took place in a time when resources were needed. This has yielded us the present day vast area of forest, still increasing in area and amount of wood. Until quite recently, forests and forest policy have been regarded as a sovereign issue in Europe, with few international measures. This diversity also led to

FIGURE 1 Forest probability map of European forests The EU27 countries cover some 178 million ha; Europe's main form of land use. Source: Gunia et al., 2011

■ Gert-Jan Nabuurs



fragmentation in research, and a concentration on national ecological and silvicultural aspects. Now, however, we stand for ever larger and more complex forest-related issues, that are cross sectoral, and require a multi-national effort. Only with this multi-national and interdisciplinary effort the required innovation in forest and forest products research can take place. Scenario studies are one way of gaining insight in alternative futures of European forests.

# Characteristics of Scenario studies in general

By 2050 the world will have changed in ways that are difficult to imagine – as difficult as it would have been at the end of the 19th century to imagine the changes of the following 100 years. Scenarios that are based on contrasting storylines can be used as a tool to explore the different ways in which the future may develop and their impacts on the sustainability of European forestry.

These scenarios are neither predictions nor forecasts, but are used to create a consistent image of a future, and to help decision makers and other actors make well informed choices (Arets et al.2008). Each story-line assumes a distinctly different direction for future developments, and does not necessarily aim to be realistic. Conclusions should not be drawn from these storylines.

A set of scenarios aims to describe divergent futures that encompass a significant portion of the underlying uncertainties in the main driving forces. These drivers cover a wide range of key characteristics such as demographic change, economic development, and technological change. For this reason, their plausibility or feasibility should not be considered solely on the basis of an extrapolation of *current* economic, technological, and social trends.

Reference futures (a baseline) and policy scenarios should be separated. Reference futures are 'benchmark' scenarios with dynamics, but without major policy interventions. Then subsequent comparison with policy scenarios enables the assessment of the effect certain policies will have.

Since driving forces can take different directions, it is better to develop multiple baseline scenarios. It is not recommended to use three alternatives because practice shows that policy makers then tend to focus on the middle scenario, which is believed to be the most realistic (Alcamo 2001).

In 1996 the IPCC decided to develop a new set of emission scenarios that are described in the Special Report on Emission Scenarios (SRES) (IPCC 1997). This set of scenarios is now known as the SRES scenarios, which were used by the IPCC for their third and fourth assessments. The scenarios are mostly developed for energy system parameters and related emissions. The underlying 4 reference futures, however, provide consistent storylines on the development of drivers like population growth and economic development in the future.

# Developments in the multi sectoral surroundings

Forests have to play their role in mitigating climate change, in providing biomass for bio energy, in pro-




FIGURE 2 Total damage due to disturbances in Europe Insight in future development of these disturbances will help decision makers make the right choices. Scenario studies can provide that insight, as well as into multi sectoral causes of these disturbances. Source: Schelhaas et al., 2008

viding jobs and opportunities in the rural areas of Europe, i.e., playing their role in the green economy. Furthermore, the vast land covered with forest needs to be managed sustainably and to protect soils and water, provide space for tourism, sequester carbon, and protect biodiversity. All this, under the impacts of climate change.

To meet these needs, changes in governance, institutions, and research structure need to take place in order to be much more innovative along the whole chain from genetics, to forest management, to harvesting, and via a variety of products, back to recycling. Also, the multi sectoral and integrated aspects need to stand out much more in research. Only then, the sustainability in all its aspects can be guaranteed.

At the international policy level, forest policy is moving to an international level in Europe. In June 2011, Ministers responsible for forests met in Oslo at the FOREST EUROPE Ministerial Conference on the Protection of Forests in Europe. They made the far-reaching decision to launch negotiations for a Legally Binding Agreement on Forests in Europe. This is a first step towards pan European forest legislation. Also the EU forest strategy is being revised. Maybe through these international policy measures, developments as the economic crisis can be turned into a break point for new developments of the forest sector.

The financial crisis which developed in autumn 2008 and the subsequent economic consequences namely have led to a fundamental shift within forestry and the forest industry sectors, the effects of which are being felt through the entire chain from forest to markets. Never, since the oil crisis in the 1970s, have the forest products markets experienced such a downturn. In response, some countries have implemented economic stimulus packages to tackle the crisis and to promote a move towards a greener economy.

Recently, attention has shifted towards economic objectives and the green economy: The wood and forestry sectors can make a significant contribution towards meeting green economy objectives, linked to climate change policies, mainly through the abatement of greenhouse gas emissions and expansion of renewable energy objectives. There are several main routes by which the wood and forestry sectors can contribute. These are sustainable wood consumption including certification and green buildings, a low carbon forest sector, creating green jobs, valuation of ecosystem services, and improved governance of the forest sector. Through this combined set of measures stimulated by government measures, and supported by research, the forest sector can really make its contribution.

## **Examples of scenario studies**

Here, we touch upon some examples of scenario studies, and the tool by which they can be carried out. Scenario studies can be carried out in a multitude of ways, but in the last decades they are more and more carried out with computerized models, aided by a fast increase in synchronised data, at ever higher level of detail. These computerized models span a wide variety, from multi sectoral global integrated models as, e.g., IMAGE (Alcamo et al. 1998), and EURURALIS (http://www.eururalis.eu, Helming et al. 2010) to specific forest sector market models and to European forest resource models. The European Forest Information SCENario model (EFISCEN; (Sallnäs 1990, schelhaas et al 2007, Nabuurs et al. 2006) is a large-scale forest scenario model that projects forest resource development on regional to European scale (Eggers et al. 2008). Here we look into three scenarios as developed for European forests (see for further details Nabuurs et al. 2006).

# Nature oriented management

It was assumed that NOM is going to be important for European forestry in the future and will reduce a forest owner's willingness to supply, i.e., as a result of less dependency of the forest owner's income on the forest, and the valuation of other functions the average owner is assumed to supply less at the same stumpage prices. This reduction in supply willingness is incor-



porated for the 27 European countries as a combination of the following assumptions:

- longer rotations (20 years for long rotations, and 10 years for short rotation species (<60 years)). This was kept rather simple because of a lack of detailed information on how the management of each tree species may change under NOM;
- from total fellings an additional 10% must originate from thinnings/group fellings;
- thinning can only be carried out in forests with growing stocks over 150 to 300 m<sup>3</sup>/ha, depending on the forest type. This is based on the assumption that non-commercial thinnings are not being practised anymore;
- a species change towards the more natural/indigenous species is incorporated as a 30 to 40% chance that species like spruce and pine will be regenerated with species like beech and oak. The accompanying assumption is that sufficient sites are available where this is a logical step;
- set aside from harvesting all beech and oak forests older than 150 years. Initially, this usually affects 1 to 1.5% of the total forest area in a country. Due to ageing of the forest during simulation, this area may increase to some 6-10% by 2060 depending on management regimes, felling levels, etc. These forest areas remain part of the simulation, but are simply not affected by harvesting anymore.

## Carbon credits

Amount of new areas being planted due to Kyoto Protocol measures

Changes in forest areas are already taking place at the moment without any carbon credits being paid. The average annual net changes in the forest area during the period 1983-1993 were highest in France and Spain with, respectively, 61.6 and 86 thousand hectares annually. Belgium, Serbia & Montenegro and Albania had seen an overall decrease in forest area<sup>2</sup>. However, these are the net changes between the gross increases and decreases in forest available for wood supply (FAWS) and forest not available for wood supply (FNAWS). For the 27 countries under study, an average annual increase in FNAWS of 324,200 ha and in FAWS of 103,600 ha has been reported. Thus, there is an overall increase in forest area of some 0.3% but only part of it is available for wood supply.

It was assumed that Article 3.3 of the Kyoto Protocol will indeed stimulate the gross FAWS area expansion: from the current +103,000 ha per year to 290,000 ha per year (on average over the whole simulation period). This scenario assumption will increase the total forest available for wood supply in the 27 European countries from the present 134 million ha (in our database) to 150 million ha. This increase was assumed to take place mainly between 2010 and 2040 and to apply to the present forest area per country with some emphasis on pre-accession countries.

Likelihood that forest owners will be financially compensated for building up carbon (growing stock) in the existing forest.

Additional carbon credits can be gained from 'forest management' up to the maximum amount individually defined for each Annex I Party (available as annex to UNFCCC, 2001). As it is in line with a strong trend in forestry, owners may be interested in it, provided that it is paid for Taking all this into consideration, as well as taking into account the high uncertainty level in outcomes of future international climate negotiations, it would be fair to assume that Article 3.4 may lead to a prolongation of rotation lengths by 10 years (irrespective of country or site). However, rotation length prolongation was mentioned under owner behaviour, NOM, and now under Kyoto Protocol issues. If forests were subjected to prolongation under all these issues, it might have resulted in assumed prolongated rotations of an extra 30 to 40 years. This seemed unrealistic and a total maximum of 20 years prolongation was assumed as a constraint.

#### Bio-energy

In the EU policy on bio-energy (EC 1997) the EU aims at doubling the contribution of Renewable Energy Sources (RES) from a current 6% to 12% by 2010. The European Commission has designated biomass as an important Renewable Energy Source on top of the current consumption of roundwood and industrial residues of approximately 40 million m³ y⁻¹ in EU and EFTA countries. In 2005, the European Commission published a Biomass Action Plan (EC, 2005b), followed by a communication on an EU Strategy for Biofuels (EC, 2006). The Biomass Action Plan aims to increase biomass use to 150 Million tonne oil equivalents (in primary energy terms) in 2010 or soon after.

Due to the above mentioned RES policy, an increase in demand for wood fibres from forest resources for the production of bio-energy has already been recorded and it can be expected to increase further. The increased demand for roundwood, based on the EU Whitepaper, has been calculated to amount to approximately 92 million m³ by 2010. Later, Lindner for EEA (2006) assessed an availability of environmentally compatible biomass from forestry of around 39 million tonne oil equivalent (~200 million m³ from both stemwood and branches).

From the current state of implementation of the RES policy in Europe, it can be concluded that it is unlikely that the RES targets for woody biomass will be met within the intended time span. Adjustment of the time span or of the quantitative targets seem inevitable. An extra felling of 80 million m³ of roundwood by 2025-2030, matches this requirement and is incorporated as an assumption in the 'new management trends' scenario. This additional fellings is distributed over the countries of study with respect to their current share in total fellings.

Combining all these trends mentioned above is called the new management trends scenario. Not applying these trends is called the 'projection of historical management' still with increased demand, stable demand is called 'benchmark'.

#### Results

Foremost, the results show that a large increase in supply can be achieved sustainably in European forests.





FIGURE 3 Actual supply under the various scenarios Source: Nabuurs et al., 2006





The supply under 'new management trends' increased from 409 million m³ y¹ in 2005 to 647 million m³ y¹ in 2060. Under the projection of historical management, the supply increased to 729 million m³ y¹ in 2060 (Figure 3). So, despite an 80 million m³ higher demand, the new management trends give a reduction in actual fellings of 82 million m³. Thus, in total a reduced supply or shortage of 162 million m³ y¹ was found despite an increase in forest area of 16 million ha over a time period of 55 years in the new management trends scenario. When keeping the felling level very stable (Benchmark), the average growing stock rises from 188 to 287 m³ ha¹. Higher rates of mortality were found under these growing stocks.

The high supply levels were found within European forests, because we allow temporarily overharvesting in comparison to increment (Figure 4). Overharvesting occurs more severely in the 'projection of historical management' scenario. Still the average growing stock increases in this scenario. In all regions where overharvesting occurs (usually starting in 2040), this happens often with an increasing linear trend in time. This in contrast to the 'new management scenario', where overharvesting is severe in some regions in 2030 and 2040, but then declines again in 2050 because of the constraints taking effect, i.e., the available stock is used up in a certain decade.

In the new management trends scenario we quantified the large scale effect of management constraints. It becomes clear that effects of management changes like setting aside forests for nature reserves, tree species changes, and rotation prolongation have their influence on total fellings mainly in the long term (Fig 3). Namely only after 2050 the fellings, as achieved under 'new management trends', start to decline. Without the afforestations the decline in fellings would have been stronger, but still the afforestations did not fully compensate for the effect of the constraints.

When keeping the felling level very stable (Benchmark), the average growing stock rises strongest from 188 to 287 m<sup>3</sup> ha<sup>-1</sup>. The age class distributions (Fig 5)




FIGURE 5 Age class distribution in 2000 and in 2060 of all European forests under simulation in the benchmark (top) and the new management trends scenario (bottom)

Source: Nabuurs et al., 2006

also show a combination of effects of different measures. Although the aim of the 'new management trends' scenario was to enhance biodiversity by creating more old forests, this is only partly achieved. We can see in Figure 5 (bottom) that the area of forests older than 200 years has increased from 1.7 million ha in 2000 to 4.4 million ha in 2060. However, the average age has hardly increased because in this scenario we also try to increase fellings a lot in combination with prolongated rotations. This leads to more forest area in 2060 in the age classes 100 to 150, but leads to less forest area in the classes 150 - 200. That is, prolongating a rotation means that you have to find the same amount of wood, but in the older age classes! Figure 5 (top) shows, for the benchmark scenario (with simply a stable and rather low amount of fellings), that the ageing of the forest is much more pronounced in this case in the classes 80-200 years.

### Concluding

Models and scenario analysis can provide a very powerful tool for explorative studies, provided they use reliable input data, and are based on thorough understanding of ecosystem functioning as well as of needs and demands from society (Mohren 2003). Large-scale scenario models in combination with land cover data

provide a means for upscaling and expansion of

it should be made very clear that results from scenario studies should not be viewed as blueprints for future developments, but rather as explorations of possible futures.

otes

1 This is not completely true anymore according to updated TBFRA data as published by UNECE/MCPFE's State of Europe's Forests, 2002

eferences

- Alcamo, J. 2001. Scenarios as tools for international environmental assessments. Experts' corner report, Prospects and Scenarios No 5. Environmental issue report No 24, European Environment Agency, Copenhagen, Denmark.
- [2] Alcamo, J., R. Leemans and E. Kreileman (eds), Global change scenarios of the 21st century. Results from the IMAGE 2.1 model. Pergamon & Elseviers Science, London. pp 296, 1998.
- [3] Arets, E., T. Palosuo, A. Moiseev, G.J. Nabuurs, D. Slimani, C. Olsmat, J. Laurijssen, B. Mason, D. McGowan, D. Vötter. 2008. D 1.4.7. Reference futures and Scenarios for the European FWC. Eforwood deliverable to Brussels. 32p.
- [4] European Commission (2005) Communication from the European Commission on the Biomass Action Plan, COM(2005)628.
- [5] European Commission (2006). Communication from the European Commission on An EU Strategy for Biofuels, COM(2006)34.
- [6] Eggers, J., M Lindner, S Zudin, S. Zaehle, and J Liski 2008. Impact of changing wood demand, climate and land use on European forest resources and carbon stocks during the 21st century. Global Change Biology 14: 1-18.
- [7] Gunia, K. Et al. 2011 www.efi.int/portal/virtual\_library/information\_services/mapping\_services/forest\_map\_of\_europe
- [8] Helming, J.F.M., Janssen, S., Meiil, van, H., en Tabeau, A., (2010).

- Impact assessment of post 2013 CAP measures on European agriculture. LEI report 2010/085. The Hague.
- [9] IPCC 2000. Special report on emission scenario. IPCC. Cambridge University Press, 599 p.
- [10] Nabuurs, G.J., J. van Brusselen, A Pussinen. M.J. Schelhaas. 2006. Future harvesting pressure on European forests. European Journal of Forest Research. 126: 391–400.
- [11] Sallnäs, O., 1990. A matrix growth model of the Swedish forest. Studia Forestalia Suecica 183. Uppsala, Swedish University of Agricultural Sciences, Faculty of Forestry, 23 pp.
- [12] Mohren, G.M.J. 2003. Large-scale scenario analysis in forest ecology and management. Forest Policy and economics. 5:103-110.
- [13] Schelhaas, M.J., J Eggers, M Lindner, G.J. Nabuurs, A Pussinen, R Paivinen, A Schuck, P.J Verkerk, D.C. van der Werf and S Zudin. 2007. Model documentation for the European Forest Information Scenario Model (EFISCEN 3.1.3). Alterra Report 1559. 118 p.
- [14] Schelhaas, M.J., 2008a. Impacts of natural disturbances on the development of European forest resources: application of model approaches from tree and stand levels to large-scale scenarios. Dissertationes Forestales 56, Alterra Scientific Contributions 23.
- [15] Von Carlowitz, 1713. Sylvicultura oeconomica.
- [16] UN ECE/ FAO 2011. The European Forest sector Outlook. Study II 2010-2030. ECE/TIM/SP8. Geneva. 111 p.